

# Atlantic salmon freshwater ALLER INFA EX GR, 0.1-0.4 MM



## Fry Feed



### DECLARATION

	0.1 mm	0.2 mm	0.4 mm
Crude protein (%)	64	64	64
Crude fat (%)	10	10	10
NFE (%)	6,9	6,9	6,9
Ash (%)	12,1	12,1	12,1
Fibre (%)	1,0	1,0	1,0
P (%)	1,4	1,4	1,4
Gross energy (MJ)	19,4	19,4	19,4
Digestible energy (MJ)	18,0	18,0	18,0

### COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

fish meal, fish oil, functional ingredients, grain products, krill meal, vegetable proteins, vitamins and minerals.

### RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

Fish (g)	Granulate	Water temperature (°C)									
			2	4	6	8	10	12	14	16	18
-	0.1 mm	0.1	0	0	0	0	0	0	0	0	0
0.2-0.3	0.2 mm	0.2	1,57	1,85	2,31	2,64	3,38	4,16	4,49	4,63	4,44
0.3-0.5	0.4 mm	0.4	1,37	1,61	2,01	2,29	2,94	3,62	3,9	4,02	3,86

### ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

	0.1 mm			0.2 mm			0.4 mm		
Feed conversion	0	0	0	0,5	0,6	0,7	0,6	0,7	0,8
N in faeces (kg)	0	0	0	0,31	0,37	0,43	0,37	0,43	0,49
N in water (kg)	0	0	0	2,06	3,03	3,99	3,03	3,99	4,95
P in faeces (kg)	0	0	0	0,23	0,27	0,32	0,27	0,32	0,36
P in water (kg)	0	0	0	0,1	0,2	0,31	0,2	0,31	0,41

### ENVIRONMENTAL IMPACT MEASURED IN CO<sub>2</sub>-EQ

Figures are in CO<sub>2</sub>-equivalents (kg/kg feed)

	0.1 mm	0.2 mm	0.4 mm
CO <sub>2</sub> -eq. with land use change	1,14-1,32	1,14-1,32	1,14-1,32
CO <sub>2</sub> -eq. without land use change	1,10-1,20	1,10-1,20	1,10-1,20