

TRADITIONAL LIPOSUCTION/FACETITE/BODYTITE LIPOSUCTION AND FACETITE/BODYTITE SKIN TIGHTENING

LOCAL ANESTHESIA

POST OPERATIVE INSTRUCTIONS

ITEMS TO PURCHASE BEFORE SURGERY:

1. **ABD pads, surgical pads, or large maxi pads.** These will help to absorb any fluid that may be seeping from the incisions (which is normal). Some patients also elect to purchase “chucks” to place on top of sheets to catch any drainage or use old sheets for the first 72 hours. Please keep in mind that drainage from incisions is typical for the first 72 hours following surgery.



2. **Compression garments:** See photos below in the “compression garment section” for examples of garments. You will need to purchase your garment prior to your surgery date. Some patients choose to purchase more than one garment. You will require these garments approximately 3 days after surgery so please plan to purchase your garments accordingly. See “compression garment” section for detailed information

3. **Lipo Foam Compression Ab Board Post Surgery Abdominal Board Liposuction (\$13):** This is to be purchased if you are ONLY having liposuction to the abdomen/flank area (not combined with a tummy tuck). The board will help ensure even pressure and improved comfort. This is especially important when sitting extended periods of time to prevent indentations. Can be purchased online at:

https://www.amazon.com/Compression-Board-Surgery-Abdominal-Liposuction/dp/B08V1KPX7P/ref=sr_1_3?crd=CYBK16TM7FKH&keywords=liposuction+foam+compression+abdominal+board&qid=1658928729&srefix=liposuction+foam+compression+abdominal+board%2Cps%2C117&sr=8-3

4. **Tight tank or kami:** If the abdominal binder is making you itchy, you may wear a tight kami shirt under your binder.

5. **Stool softener/laxative (Peri-Colace)**—Pain medication can constipate you. Peri-Colace helps soften the feces (“the mush”) and also helps stimulate the bowels (“the push”). These can be started 2 days prior to surgery to help “prep” the bowels and should be continued after surgery as long as you are taking narcotics.

6. **Extra-Strength Tylenol (500mg):** To be used when pain is mild to moderate in place of the narcotic pain medication.

7. **Hibiclens**® antibacterial liquid: You will use this liquid the morning of surgery to wash the surgical areas. You do not need to use this soap in the postoperative period as it will dry out your skin. You may resume your normal soap postoperatively on day 3 after surgery.

8. **Arnika and Bromelin**: These are herbal supplements that aid in bruising and swelling. This supplement is included in your surgical cost and will be provided at your pre-operative appointment. You can begin these supplements 2-3 days before surgery and use them for 2 weeks after surgery.

9. **Compression stockings**: We will provide you with 1 pair of compression stockings immediately after surgery. Please plan to purchase an additional pair at Target® if your socks become soiled or are uncomfortable. They will need to be worn for 5 days postoperatively.

10. **Oral Thermometer**: It is important to monitor your temperature after surgery if you feel “feverish” or hot. You have a fever if you have a temperature above 101.5.

11. **Comfortable Clothing**: Please wear comfortable clothes to your surgery. Loose-fitting, button-down shirts and/or maxi dresses work well.

WHAT TO EXPECT AFTER SURGERY

- Abdominal bloating is to be expected in the early postoperative period (even if this is not the area of treatment). Bloating results from sodium present in your medications and in the anesthesia.
- Most patients report difficulty sleeping and difficulty with pain management the first night after surgery. This is usually secondary to the effects of anesthesia and should subside within 24-48 hours.
- Moderate swelling of areas of treatment is to be expected. You may find that your clothes do not fit as easily as before. Be patient. The swelling will gradually subside, and you will be back to normal in 3 – 6 months. Swelling will be at its worst between 3 – 5 days. Swelling starts to subside at 6 weeks but may take 6 months to resolve.
- If the abdomen/flank area is treated with liposuction; swelling to the hips is common for the first 1-2 weeks. This is due to the abdominal binder used after surgery. The swelling will improve once you switch to your full body compression garment.
- You will not see immediate weight loss for the first 2-3 weeks. In the long run, you will see more of a change in how your clothes fit than a decrease in weight.
- To alleviate any discomfort and reduce swelling, you may apply cool, not cold compresses to the treated area. Frozen bags of peas and corn work well; make sure to wrap in a towel before applying to the skin. Apply cool compress for no longer than 10 minutes at a time.
- Bruising is a normal expectation following surgery. Bruising could be apparent for as long as 3 – 4 weeks afterwards. The bruises will move down your body as they are absorbed and will change in color from a dark purple to yellow as it clears.
- A burning sensation (raw, sensitive) or sharp shooting pains along the surgical area and incision line is normal and indicative of nerve regeneration. These “neuralgias” will burn out by 9 months.
- Numbness is expected to resolve by 9 months on average.

- Shiny, itchy skin due to swelling. If this occurs, an OTC antihistamine such as Benadryl can help alleviate symptoms.

- It takes **6 MONTHS FOR FINAL RESULTS** to appear. In the interim, you may notice:

- Incision asymmetry
- Surgical site asymmetry
- Swelling changes from day-to-day.
- Redness to the incisions that may take up to 9 months to fade.
- Be patient please and try not to focus on these issues before the 6-month period.

- ***Consistent, sharp pain that is unrelieved by pain medication and/or repositioning should be reported to our office immediately.***

PLEASE CALL THE OFFICE AT (972) 930-0333 IF:

- You have redness, increased pain at surgical incision sites, sudden increase in swelling, warmth, drainage (pus), or oral temperature greater than 101.5°F

PRE/POST OPERATIVE MEDICATIONS

1. **Norco** (pain medication): see below.
2. **Valium** 5mg 1 tab po morning of surgery and 12 hours later (if needed).
3. **Zofran** (nausea) 4mg 1 tab every six to eight hours as needed for nausea. Start after discharge.

****Please bring all medications the morning of surgery. The OR team will dispense the medication upon your arrival****

4. **Silagen** Arnica -Bromealin Supplements:
 - a. A unique formulation that is clinically proven to speed the healing time up to 50% allowing for faster recovery from bruising, swelling, and pain associated with surgical procedures.
 - b. Start 2-3 days prior to surgery with 1 tab in the morning and 1 tab at night. You can continue this after surgery when you are actively swollen and bruised.



Notes on Pain Medication:

- Take pain medication with food.
- Norco (Hydrocodone + Acetaminophen) should be taken as directed:

- Mild to moderate pain: 1 tablet every 4 – 6 hours as needed
- Severe pain: 2 tablets every 4 – 6 hours as needed.
- If your pain is mild, or if you do not like the effects of the narcotics, you can take Tylenol® (Acetaminophen) 1000mg every 8 hours (which would be 2 tablets of the Extra Strength Tylenol®, purchased over the counter). Please do not exceed 3000mg in a 24-hour period. Please do not mix the Tylenol® with the narcotic pain medication since the narcotic pain medication consists of 325mg of Tylenol®
- We will provide every patient with 1 refill of the pain medication if needed one week after the last prescription was filled. No additional refills will be provided.
- If you are to finish your pain medication during a weekend or after hours, you will need to wait until regular business hours to request additional pain medication.
- If you are under the care of a pain management provider or already take narcotic pain medication, our office **WILL NOT** be providing you with an initial RX or refill for the aforementioned. It is your responsibility to obtain the necessary medications/refills from your prescribing provider. Please make sure to bring the necessary medications if you are visiting us from out-of-town as we will not be providing it.
- Our office will not be providing any additional medications that are unrelated to the surgical site (i.e., blood pressure meds, muscle relaxers, etc.). Should you require these in the postoperative period, you will be responsible for acquiring them from your primary care provider.
- Take a stool softener with pain medication to prevent constipation.
- **DO NOT DRIVE WHILE TAKING PAIN MEDICATION.** These medications can result in drowsiness. If you are pulled over while driving under the influence of narcotics or scheduled substances, you will get a DUI!
- **DO NOT DRINK ALCOHOL WHILE TAKING PAIN MEDICATIONS.** This can be a deadly combination.
- Only take the narcotic pain medication if needed. The quicker you can wean off the pain medication, the better you will feel and heal.

HAVE SOMEONE WITH YOU

- After surgery, have an adult available to drive you home and stay with you for the first 24 hours, as you will be weak and drowsy.
- It is highly recommended that you have an adult with you for the duration of your stay if you are an out-of-town patient.

DO NOT SMOKE. This is very important!!!

- Smoking (tobacco, marijuana, or vapes) can result in a lack of blood supply to tissues and fat causing tissue death or delayed wound healing. Even 0% nicotine vapes contain a trace amount of nicotine that the FDA accepts as 0%.

- Do not allow those caring for you to smoke around you as second-hand smoke can be detrimental to your recovery.
- No need to resume smoking as you have stopped six weeks before surgery. This is a benefit to your overall health.

SUPPLEMENTS

- Do not take aspirin (or products containing aspirin), anti-inflammatories, or Ibuprofen (Advil®, Motrin®, Midol®) for 1 week after surgery. Also do not begin herbal supplements until 1 week after surgery.

- Arnika, Bromelin and Vitamin C are okay to take.

- Phentermine or appetite suppressants should not be taken until 6 weeks after surgery as these supplements increase heart rate and blood pressure and can interfere with your recovery.

WALKING

- It is important to get out of bed early and often after your surgery (with assistance) to prevent postoperative problems. Walking encourages blood flow throughout your legs to reduce the chance of blood clot development.
- IF YOU HAVE SHORTNESS OF BREATH, LEG SWELLING, AND/OR LEG PAIN AT ANY POINT IN YOUR POSTOPERATIVE HEALING, GO TO AN EMERGENCY DEPARTMENT IMMEDIATELY (OR CALL 911) AS THIS COULD SIGNIFY A BLOOD CLOT.

SLEEP POSITION:

- While resting in bed, keep a pillow under your legs and at least 2-3 pillows behind your back and head. This position will help aid in the reduction of swelling.
- While sleeping, sleep at approximately 30—45-degree angle. Again, this is best achieved with pillows behind the head/back and knees. Best to remember, whatever area has been treated, try to elevate that surgical area.
- After two weeks you can slowly start to lay flat on your back. Side sleeping can be achieved closer to the 3-4 week post operative period.

SHOWERING

- You may shower (but not bathe) 48 hours after surgery.
- Make sure someone is with you at your first shower. You can expect some dizziness with your first shower. Make the shower a quick one.
- You may wash the surgical area with a mild soap and water (lukewarm, never hot). Do not use surgical soaps to wash the area as these are drying to the skin. Use regular soap.
- Remove all your garments, dressings when showering. When in the shower, use your hands to gently wash the surgical sites to remove dry blood, sweat and oil. When out of the shower, pat the incisions

dry. There is no need to use dressing covers if you are not draining from your incisions. The incisions may encounter the compression garments.

- Moisturize the surrounding skin around the incision line if the skin becomes too dry and/or irritated.

DIET

- Stay on a soft diet, high in protein, for 2 – 3 days and avoid spicy food which can cause nausea and gas. Then you may resume a normal, high protein diet.

EXERCISE AND SEXUAL ACTIVITY

- No overheating for the first 3 weeks (spas, exercising in the sun, etc).
- Elevating the heart rate, i.e., brisk walk/treadmill may begin starting at 2 weeks.
- At 2 weeks you can consider passive, or less aggressive sexual activity.
- At 3-4 weeks, you can consider slowly reintroducing your workout routine.
- You may begin swimming 3-4 weeks post-operatively if healing is complete. We advise to wait a full 6 weeks before swimming in a lake or ocean.

SUTURES

- You will have non-absorbable sutures in place. Sutures will be removed at your one week post operative appointment.
- Do not apply anything on your incisions for 3 weeks unless told to do so. Keep the incisions clean and dry.
- At 3 weeks post operatively, you may begin scar treatment with Silagen®.

SCAR THERAPY

- We use and sell Silagen® Scar Refinement System at our office. This system will be discussed at the three week post operative appointment. Please note that you will receive one pump at your pre-operative appointment. You can purchase additional Silagen® at our office.
 - Silagen® is made with the highest quality medical grade silicones that create a protective barrier over scars which increase hydration and help stop excessive collagen buildup. This will help flatten and soften scars and reduce redness, itching, and pain.
- You may begin using Silagen® as soon as the skin is fully closed, after all sutures are removed and after all scabs have fallen off. This usually occurs around 3 weeks from surgery.
 - We recommend you use at Silagen® daily for approximately 6 months as that is the time you can achieve maximum benefits and results. You can continue to use Silagen® up until the one-year post surgery date.




- All incisions will be extremely sensitive to sunlight during the healing phase. Direct sun contact or tanning booths are to be avoided for 9-12 months. Use a water-resistant sunscreen with SPF of 50+ with UVA and UVB protection for at least 9-12 months. Sun damage to the scars may result in permanent hyperpigmentation or hypopigmentation to your scars.
- Please note that use of medical grade silicone sheeting or silicone cream on surgical scars will temporarily result in red/purple pigment on the scars. This takes place as blood rushes to the surgical incision to create collagen for wound healing and scar maturation. It can take 9 months on average for the discoloration to begin to fade.

COMPRESSION GARMENTS

- Wear the white abdominal binder (compression garment) that we provide for approximately 24-48 hours. *This applies only if having liposuction to the abdomen and/or flanks.*
 - The white abdominal binder will be in place until all the drainage/leakage from the liposuction areas has ended.
 - Often patients like to wear a tight shirt or cami under the abdominal provider to help alleviate any itching the binder causes.
- Once the drainage/leakage has stopped, you are able to wear the full body garment you ordered prior to surgery.
- Where to buy:
 - Our office likes to utilize Rashida Carter, a specialist in compression garments and lymphatic massages. She wholesales medical grade, Colombian fajas. Rashida will measure you prior to surgery and guide you on purchasing the appropriate garment.
 - Rashida Carter
 - 10300 North Central Expressway Suite 461, Dallas, TX 75231
 - (469) 264-9786
 - Business card will be provided in pre-operative appointment.
 - Additional garments can be purchased, however; our advice is to wait until the swelling has decreased as you may require a tighter fitting garment, than what is used right after surgery.
 - If you elect not to use Rashida's services, you can purchase compression garments on sites such as Marena, Amazon, etc.
- Garment time frames:
 - **Body** (not including arms): garment will be worn 24/7 for the first 6 weeks following surgery.
 - **Arms:** garment will be worn 24/7 for the first 3 weeks following surgery.
 - **Face:** garment will be worn 24/7 for the first 3 days and then only in the evenings for 2-3 weeks.

- Please bring your face garment the day of surgery. The OR team will put the garment on you after surgery.



COMMON COMPLICATIONS

• SEROMA

- Some patients may experience fluid build-up in the areas of treatment in the early postoperative period. If this happens, you will notice a fullness or sloshy feeling in the specific area.
- If a seroma is detectable (palpable or visible) then it can be treated through aspiration. A small needle and syringe will be used to drain the fluid out of the surgical site through your incision which is typically numb.
- A patient with a seroma may require serial weekly aspirations. Typically, as a patient's body heals, less fluid will be recruited into the area. A seroma will resolve within 3 - 6 weeks with appropriate treatment.



• KELOIDS/HYPERTROPIC SCARS

- Keloids are firm, rubbery, lumpy lesions or shiny fibrous nodules on the skin that's usually raised. When skin is injured, fibrous tissue called scar tissue forms over a wound to repair and protect the injury. In some cases, scar tissue grows excessively, forming smooth, hard growths called keloids.
- Hypertrophic scars are raised scars that do not grow excessively outside the incision line.

- If keloids or hypertrophic scars are detected, treatment may include:
 - Inject corticosteroids to the area to reduce inflammation (usually one injection every 6 weeks for up to four sessions)
 - Continue silicone gel sheeting
 - Begin laser treatments and micro needling to reduce scar tissue and help with discoloration.



INFECTION

- Infections are most likely to take place between day 10-14 from surgery.
 - Symptoms and signs of infection include redness, warmth, fever, tenderness, pus, malaise
- Most infections, if detected early, can be treated with proper antibiotic therapy. If infections are severe or not responding to antibiotic therapy, then prompt visit to an ER for IV antibiotic will be necessary.
- If you are an out-of-town patient and you develop an infection, you will need to visit with your local provider or local ER for evaluation and care. If fever and pus accompany redness, please visit with your local ER.
 - It is not recommended that you travel long distances with an infection.

PLEASE VISIT AN EMERGENCY ROOM OR CALL 911 IF:

- At any point you experience shortness of breath or leg pain with swelling as this could indicate a pulmonary embolism (blood clot in lung) or DVT (blood clot in legs) and could be deadly if untreated.

FOLLOW UP APPOINTMENTS

- It is important to be seen by our office for your post-op checks.
- Dr. Setty will see you at the following follow-up appointments: one week, six weeks, six months, and one year.
 - 1 Week Post Operative Appointment
 - Dressing change (if needed)
 - JP drain removal (if needed and if under the 30cc amount in a 24-hour period)
 - Suture removal
 - Observe tissue for necrosis or wound separation
 - 6 Week Post Operative Appointment
 - Assessment for scar management (i.e., if steroid injections, topical creams, and/or laser treatments are needed.
 - Post operative photos started.
 - 6 Month Post Operative Appointment

- First conversation regarding aesthetic results/concerns.
- Post operative photos.
- One Year Post Operative Appointment
 - Final conversation regarding aesthetic results/concerns.
 - Remember your body will change with age which also means the appearance of your treatment areas will change too. Although the outcomes of a liposuction are generally permanent, any significant weight gain or loss, as well as the normal influences of aging can cause changes to your appearance. You may wish to undergo revision surgery later to help maintain your appearance throughout life. Contact our office with any of your questions or concerns, at any time.
- RN will see you at the following appointments: three-week, three-month, six months.
 - 3 Week Post Operative Appointment
 - Assessment of surgical site.
 - Initiate treatment with scar therapy.
 - 3 month Post Operative Appointment
 - Measurements and post operative photos.
- *It is strongly advised that you stay locally for the first week following surgery if you are an out-of-town patient.*
- It's important to keep in mind that these appointments are patient-specific and may vary depending on your own individual healing and/or complications.
- If you are from out-of-town and cannot attend appointments at the recommended frequency, then it is strongly advised that you establish a relationship with a provider in your hometown who can follow your progress or evaluate you in case of infection, wound separation, or seroma. Virtual appointments may be scheduled if unable to attend in-person appointments.
- Follow up appointments can be made while at your scheduled post operative appointment or you can call the office at (972) 930-0333. Our office hours are Monday – Thursday 8:00 AM - 4:00 PM and Friday 8:00 AM – 3:00 PM. The office is closed on Saturday and Sunday.
- **The office is closed on Saturday and Sunday. Should you experience a complication over the weekend, call the regular office line at (972) 930-0333 and you will be directed to an afterhours line or if it is a life-threatening emergency, please visit the nearest urgent care/ER.**

ITEMS TO CONSIDER TO AIDE IN RECOVERY

1. **Lymphatic Drainage Massage:** Consider this therapy to decrease healing time up to 50% by massage techniques used to reduce swelling and pain after surgical procedures. It's also known to reduce overall recovery time and allows patients to be more comfortable during the recovery process. Light pressure and strokes are used to encourage the flow of lymph material to be filtered out of the body. Referral information will be provided in your pre-operative folder.

2. **Hyperbaric Oxygen Therapy (HBOT):** Consider this therapy to decrease healing time up to 50% by reducing reperfusion injury, swelling and inflammation. HBOT also improves blood circulation, reduces pain, and upregulates collagen production.

3. **Wound Care Center and Primary Care Provider:** (for out-of-town patients only): If you are an out-of-town patient, please arrange care with a local wound care center and with your primary care provider. Brachioplasty can result in areas of delayed wound healing that may require additional wound care postoperatively. Should you suffer a complication (i.e., infection or fluid accumulation), you will need to be evaluated and treated by your primary care provider. Should you suffer a complication like delayed wound healing or wound separation, you will need to be evaluated and treated by your local wound care center.

IMPORTANT POINTS OFTEN OVERLOOKED ABOUT SURGERY

- It is not unusual for patients to undergo significant emotional “ups and downs” after any type of surgery. Factors such as underlying stress, medications, and/or psychological tendencies can result in patients experiencing a “post operative depression” that generally resolves after a few weeks. Having a partner, family member, or friend who is supportive can help with this process. Understanding the stages of emotional “ups and downs” can help patients stay calm and recover from this emotional process faster:

- Phase 1: Being Out of It
 - Swelling and discomfort is most severe over the first few days after surgery. Pain medications also can make you disoriented and emotional.
- Phase 2: Mood Swings
 - Having just had surgery, patients are adjusting to a sudden change in their appearance with much anticipation. The presence of bruising, swelling, and asymmetries will distort a patient’s results thereby concealing the final outcome. Mood swings (especially sadness), worry and depression are common emotions as a result. Patients may even ask, "What have I done?" or think that "I never should have done it."
- Phase 3: Being over critical
 - During the second week, patients will probably be feeling a lot better. The swelling and muscle cramping/spasms will be decreasing, and sutures will be out. Because of anticipation, it is natural for patients to look critically at their new body worrying about symmetry, scars, and so on. At this point, it's normal to wonder if they have achieved their goal and what they paid for. This is too soon to tell, and most concerns are resolved with time.
- Phase 4: Happy at last
 - Finally, about 3 – 6 months out of surgery, patients will probably start liking how they look and are feeling much better. They may be in the mood to check out some bathing suits or outfits to show off their new figure.