Patient Education

Pain Medication Information

Upon arrival for your procedure, please update your healthcare provider with the following information:

- Pain medications that you are NOT able to take for any reason, such as allergy or intolerance, OR those that do not work for you.
- Pain medications that you have used in the past that **do** work well.
- Whether you are presently taking any other pain medications.

Over-the-counter pain medication options:

NSAIDS non-steroidal anti-inflammatory drugs, include Ibuprofen, Motrin & Advil.

How they work: Nonsteroidal anti-inflammatory drugs reduce pain, decrease fever, and, in higher doses, decrease inflammation. Use with caution, and let your doctor know if you have a history of <u>kidney</u> problems. You can take **up to** 600 mg of Ibuprofen every six (6) hours as needed for pain.

Acetaminophen Tylenol. Regular strength tablets are 325 mg, extra strength are 500 mg.

How it works: Acetaminophen is a pain reliever and a fever reducer. It is used to treat conditions such as headache, muscle aches, arthritis, backache, toothaches, colds and fevers. Use with caution, and let your doctor know if you have a history of <u>liver</u> problems. You can take **up to** 1,000 mg of Acetaminophen (Tylenol) every six (6) hours as needed for pain, a maximum of 4,000 mg in 24 hours.

For best pain coverage, it is recommended that you alternate these medications every 3 hours. For example:

8 a.m. Take Ibuprofen.

11 a.m. Take Tylenol.

2 p.m. Take Ibuprofen.

5 p.m. Take Tylenol.

These are the maximum doses that can be taken. If your pain is under control, we encourage you to increase the length of time in between doses.

Page 1 of 2



Hydrocodone, Oxycodone and Tramadol are common prescription-only narcotic pain medications.

- Hydrocodone/acetaminophen 5/325 mg (Vicodin) is 5 mg of a narcotic (hydrocodone) mixed with 325 mg of Tylenol.
- Oxycodone/acetaminophen 5/325 mg (Percocet) is 5 mg of a narcotic (oxycodone) mixed with 325 mg of Tylenol.

IMPORTANT: If you are taking either one of these medications and are trying to transition to over-the-counter pain medications, be sure that you count the amount of acetaminophen in each of these tablets towards your recommended maximum dosage.