# **Glasgold Group Surgery Center**

## POST-OPERATIVE INSTRUCTIONS BLEPHAROPLASTY

To help with swelling, apply ice/cold compresses to your eyes for 3 days. Wrap the ice packs in clean gauze. **Do not** apply ice directly to the skin. Apply for 10-15 minutes at a time. **Do not** ice for longer than 15 minutes at a time. Apply ice on and off throughout the day as directed 10-15 minutes on, 20-30 minutes off. You may ice past 4 days for comfort.

Keep head elevated on 2-3 pillows while resting and sleeping for at least the first 48 hours.

You may shower briefly (not too hot) 24 hours after the surgery.

The 1st post-operative appointment is 6-7 days after surgery. Sutures will be removed.

**AQUAPHOR/VASELINE** ointment should be applied to the suture lines 2 times a day until sutures are removed. There is a running suture under the skin with the ends of the sutures taped at the eyebrow with a steri-strip.

**<u>Do not</u>** remove the tape or cut the suture. If the tape comes off, cut the sticky end of a bandaid and tape the suture back up.

**STERILE eye wash** solution may be used to moisten the eyes as needed and may be obtained from your pharmacy: (i.e., rewetting drops, sterile saline solution, or Natural Tears).

Extra-strength Tylenol may be taken for discomfort, or you may take the prescription pain reliever as directed instead if needed.

EYE MAKE-UP may be applied 3 days after sutures are removed. Be gentle with application and removal.

#### Additional Instructions for Patients Who Had Lower Lid Transconjunctival Blepharoplasty

Do not wear contact lenses for at least 1 week.

### **Contact Information**

During regular business hours call (732)846-6540

Before/after regular business hours call (732)846-6540 press #4

#### In case of emergency call 911 or go directly to the ER

S DRIVE/FORMS CLINICAL/POST-OPERATIVE INSTRUCTIONS BLEPHAROPLASTY as of Nov 2017