


Annual Report Brief PATIENT OUTCOMES

Ideal Option provides low-barrier treatment to people struggling with substance use disorder.



Ideal Option's Treatment Works.

Lab testing results for patients who remained in Ideal Option's treatment program and stayed on their prescribed medications show:

Treatment Works²

less fentanvl use




methamphetamine use

Life Improves³

improvement in housing



improvement in employment



Self-reported outcomes gathered via survey of over 1,800 Ideal Option patients after a minimum of 6 months of treatment show:



96% Report their quality of life has improved



93% Report no visits to the **ER** for drug-related medical care since starting treatment



96% Report no drugrelated arrests or **charges** since starting treatment



98% Report they feel confident in their ability to remain in recovery long-term



patients who returned for a second visit remained in treatment after 6 months

Source: Ideal Option 2023 Annual Patient Outcomes Report

³2023 Ideal Option Self-Reported Patient Outcomes Survey

INITIATION

New patients, re-start patients, or patients not yet stabilized on buprenorphine.



STABILIZATION A

Patients who are starting to feel stable on buprenorphine and have reduced their use of opioids.



Stages of Care STABILIZATION B

Patients who are stable on buprenorphine and have stopped use of opioids but may still be using other nonprescribed substances.

1 Visit Every

MAINTENANCE A

Patients who are completely stable on buprenorphine and abstinent from all nonprescribed substances.

L Visit Every

Maintenance A

MAINTENANCE B

Patients who have long-term stability on buprenorphine and are abstinent from all nonprescribed substances.



No Withdrawal Long-term Stability **Symptoms** No Withdrawal No Withdrawal No Withdrawal GRADUATION Symptoms **Symptoms** No Cravings Symptoms **CRITERIA** Reduced Cravings No Cravings Taking Buprenorphine* No Cravings *Verified by Taking Buprenorphine* Taking Buprenorphine* No Opioid Use* Taking Buprenorphine* diagnostic test Reduced Opioid Use* No Opioid Use* No Opioid Use* No Non-Prescribed Benzos or Sedatives* No Non-Prescribed No Non-Prescribed Benzos or Sedatives* No Stimulants* Benzos or Sedatives* Note: Patients with isolated methamphetamine or alcohol use disorder will follow specialized protocols No Stimulants*

Average Positive Rates by Stage of Care

