The drugs listed below have undesirable side effects and/or may affect your anesthesia or surgery. These drugs should be avoided for two weeks prior to surgery, and two weeks after surgery. If you need to take something for minor aches and pains, you may take Tylenol or another acetaminophen product. **NO ASPIRIN/ASPIRIN-BASED PRODUCTS OR BIRTH CONTROL PILLS/ INJECTIONS* TWO WEEKS BEFORE AND AFTER.** Aspirin will cause excessive bleeding. Birth control will increase chances of blood clots. **ALSO, TWO WEEKS PRIOR, NO **HERBAL SUPPLEMENTS, VITAMIN E or RED WINE. TESTOSTERONE MUST BE DISCONTINUED 2 WKS BEFORE, BUT MAY BE RESUMED AFTER ONE WEEK**. Please let Dr. Chegireddy know if you are currently taking any of these medications on a regular basis.

Advil
Aleve
Alka Seltzer
Anacin
Anaprox
Anadynos
Ansaid
A.P.C.
Argesic
Arthropan liquid
Arthritis Pain Formula
Arthritis Strength
Bufferin
A.S.A.
A.S.A. Enseals
Ascriptin
Asperbuf
Aspergum
Aspirin
Altromid
Axotal
Azolid
Bayer Aspirin
B.C. Tablets and powder
Birth Control Pills
Birth Control Injection
Buff A Comp
Bufferin Arthritis
Bufferin
Bufferin Extra Strength
Buffet II
Buf-Tabs
Butazolidin
Cama Arthritis Reliever
Cheracol Capsules
Chlortrimeton
Clinoril
Congesprin Chewables
Cosprin
CP-2 Tabs
Damason P
Daryon Compound
Darvon
Darvon N, with A.S.A
Dulvules
Di-gesic
Disalcid
Dolobid
Doprn
Dristan
Durasal Tablets
Easprin
Ecotrin
Efficin
Elavil
Emagrin
Empirin with Codeine
Emprazil
Encaprin
Endep
Equagesic Tablets
Estrogen Blockers
Etrafon
Excedrin
Feldene
Fiorinal
Fish Oil
Flagyl
Flexeril
Four Way Cold Tabs
Gaysal S
Ibuprofen
Imitrex
Indocin
Indomethacin
Lanorinal
Lioresal
Lortab
Magan
Magsal
Measurin
Micrainin
Midol
Mobigestic
Morbidin
Motrin
Mysteelin
Nalfon
Naprosyn
Naproxen
Neocylate
Nicobid
Norgesic
Norgesic Forte
Nuprin
Oraflex
Orudis
Osteo Bi-Flex
Pamelor
Pepto Bismol
Percodan
Perocan
Persistin
Phentermine
Phenylbutazone
Ponseil
Progesterone
Propxyphene
Compound 45
Rufen
Ru-Tuss
Sine-Aid
Sinutab
SK-55 Compound
Stendin
Tagamet
Tamoxifen
Testosterone
Tetracycline
Tolectin
Triaminicin
Trigestic
Uracel
Vanquish
Verin
Vibramycin
Vitamin E
Voltaren
Zomax
Zorprin

Please use this list as a guideline. Report the use of all medications to Dr. Chegireddy (404) 446-0044.

*Birth control injections cannot be administered within two weeks of surgery (2 weeks before and after).

**Herbal supplements are not regulated by the FDA, therefore we do not know for certain what the supplements contain. In order to ensure patient safety, we advise stopping all herbal supplements two weeks prior to your surgical procedure.