



PHONE: **304-598-4835 FAX: 304-974-3250 PO Box 9214 Morgantown, WV 26506-9214**

Name:				DOB:	/	/
School fax:		School:		Date:	/	/
School Nurse E	E-Mail:					
		chool year unless otherwis complete 504 Plan due to p				
PLEASE COM	PLETE THE FOLL	OWING:				
DIAGNOSIS:	☐ Type 1	Type 2				
THERAPY:	☐ Injection (MDI	☐ Insulin pump	□cgм			
☐ Rapid actin	g insulin includin	g Admelog, Apidra, Huma	alog, NovoLog, Fiasp	, Lyumjev		
hyperglycemia. sticks. If using Before brea Due to compating feet of the stude purposes while BLOOD GLUCO	Students may us CGM, have studer kfast Before bility of CGMs and lent at all times to prin class. DSE TESTING (Sc	nsulin pumps, all cellular de event data and communica nool nurse must determine	in dosing calculation lues on pump before seed: evices, readers, and/or tion loss. Students shows whether it is safe and	controllers should only be used appropriate):	blood sug build be ke	gar finger
☐ Independen	t ∐ Depen	dent with nurse supervis	ion or school nurse (delegation		
		FOR MEALS AND SNAC	_			
BREAKFAST:	Insulin/carbohydr	ate ratio:				
LUNCH:	Insulin/carbohydr	ate ratio:				
SNACKS:	•	ate ratio:				
SLIDING SCAL	E:					

3. SNACKS:

*Boluses may be given pre-meal or post-meal per parent discretion.

Snacks to cover low sugars do not require insulin coverage. Snacks are not required but can be given as needed. Follow the above carbohydrate order if not used for treating lows.

4. HYPOGLYCEMIA: (Signs of hypoglycemia include hunger, feeling funny, shakiness, irritable, tired, drowsy, weakness, paleness, dizziness, sweating, inability to concentrate, personality changes. If present, check blood sugar or CGM reading. Parents may modify carbohydrate grams for treating lows)

For blood glucose less than 80, use the 15/15 treatment method: Escort to nurse's office and treat with 15 grams of fast- acting carbs. Recheck blood glucose in 15 minutes and treat again if needed. If the next meal or snack is greater than 45 minutes away, please follow with a small snack containing complex carbohydrate and protein, such as cheese crackers. If low occurs right before lunch, follow the 15/15 treatment method and allow the student to eat lunch. Then, cover the carb intake subtracting 15 grams. If blood glucose is <60 when first checked, treat with 30 grams of a simple carbohydrate. Then, cover lunch carb intake subtracting 30 grams.



fluid per hour.

PHONE: **304-598-4835 FAX: 304-974-3250 PO Box 9214 Morgantown, WV 26506-9214**

 5. GLUCAGON: May be administered as needed for severe hypoglycemia. Give intramuscularly or subcutaneously if unconscious, seizure, or unable to swallow. Call 911. (Glucagon kit/Gvoke dose; Baqsimi = 3mg only) ☐ 0.5 MG OR Baqsimi ☐ 1.0 MG OR Baqsimi
Possible side effects of Glucagon include nausea or vomiting. Position student on side after glucagon administration until fully awake. Glucagon may not be self administered. Glucagon/Baqsimi or glucose gel (and other needed diabetes supplies) must be taken on field trips.
6. HYPERGLYCEMIA : (Signs of hyperglycemia include thirst, frequent urination, fatigue or sleepiness, blurred vision, difficulty concentrating, flushing of skin, sweet or fruity breath, nausea, and vomiting. If present, check blood sugar or CGM reading)
School nurse may supervise and/or administer pump bolus or injection of rapid-acting insulin for blood glucose per sliding scale. Only repeat after two hours from the first correction dose. Encourage oral hydration for BG/CGM >350 (16-20 ounces).
INSULIN ADMINISTRATION (Check one box only): ☐ Independent (without help or supervision) ☐ Dependent
☐ Independent with parent indirect supervision. Insulin administration may be progressed to independent if school nurse and parents determine this to be safe and appropriate. This is a mutual plan established by parents and school. Prior approval must be obtained.
Students should receive additional care at home, at a hospital, or through follow-up with a physician if: • Persistent hyperglycemia (blood glucose >400) after 3-4 hours of treatment and monitoring despite 2+ insulin coverage for high. Recheck blood sugar no sooner than a half hour after correction dose is given. • If patient is vomiting.
INSULIN SIDE EFFECT: hypoglycemia. Do not correct high blood sugars more frequently than every two hours with short-acting insulin.
FOR PUMP USER ONLY: Allow pump correction features to determine dosage with the input of blood glucose, meal announcement, and/or carbohydrate intake. In the event of pump failure, use the sliding scale every two hours and revert back to multi dose injections.
Any time blood sugars do not respond to correction bolus, check infusion setting and/or pumpsite. It may need to be changed, and insulin managed by injection based on the student's sliding scale. Please contact parents if this occurs.
7. URINE DIPS: Please check urine ketones if blood sugar is over 300 for four consecutive hours and/or if student is vomiting. Parents will supply ketone strips. Please refrain student from exercise when ketones are present. If positive,

Americans with Disabilities Act. Examples of such occurrences include physician appointments, hospitalization, educational classes related to diagnosis, diabetic ketoacidosis, hypoglycemia, and profound hyperglycemia. **Recurrent school absenteeism from diabetes should be rare and will not be excused by this service**.

8. ABSENTEEISM: Student is exempt from the policy if the reason is related to their diagnosis according to the

please give coverage based on sliding scale and encourage student to drink one-to-two cups of water or other sugarfree

9. FIELD TRIPS: Student may participate in field trips. Glucagon/Baqsimi or glucose gel must be taken on field trips.



PHONE: **304-598-4835**

FAX: **304-974-3250 PO Box 9214**

Morgantown, WV 26506-9214

10. FREQUENT BATHROOM BREAKS and drinks are permitted at any time not just with hypo- and hyperglycemic episodes. There should be no punitive action taken.

- 11. ACADEMIC TESTING: If student is experiencing a hypoglycemia (<80) or hyperglycemic (>350) episode prior to or during a test, the student must retake the test at a later time. Uncontrolled blood sugars can affect the thought process and ultimately affect grades.
- **12.** Please call **304-598-4835** if there are any questions or concerns.

I GIVE MY PERMISSION FOR MY CHILD TO BE TREATED PER ABOVE ORDERS AND FOR THE EXCHANGE OF PERTINENT INFORMATION BETWEEN THE SCHOOL AND DOCTOR'S OFFICE. I UNDERSTAND IT IS THE PARENT/ GUARDIAN'S RESPONSIBILITY TO PROVIDE INSULIN AND ALL TESTING AND TREATMENT SUPPLIES.

Signature of Parent/Guardian	Date:	/	/	
	Date:	/	/	

Evan A. Jones, MD, PhD Brian Ely, MD Elizabeth Minchau, DNP, APRN-BC Kathleen Sorkin, APRN, CDCES



Department of Pediatrics Pediatric Endocrinology 1 Medical Center Drive Morgantown, WV 26506-9214

10 Medical Park Wheeling, WV 26003-6389



PHONE: **304-598-4835**

FAX: **304-974-3250 PO Box 9214**

Morgantown, WV 26506-9214

INSTRUCTIONS FOR COMPLETION OF DIABETES SCHOOL ORDERS

Patients and families.

Our team places the highest priority on meeting the diabetes needs of your child as we transition into the upcoming school year. To ease this transition, we are asking that you please complete the following steps so that your child will have optimal diabetes management at school.

- Please contact your school nurse and schedule a meeting prior to the start of school.
- 2. Complete this Diabetes School Orders in its entirety prior to meeting with your school nurse. During your meeting, please discuss the information on the form and any concerns for your child's care while they attend school. Please agree upon a plan of care for your child during school hours.
- 3. Send the completed form through any of the options listed below:
 - Fax to 304-974-3250. Your school nurse may also be able to assist in faxing your forms during the meeting.
 - Send through MyWVUChart as an attachment. If you have not signed up for MyWVUChart, our office can assist you.
 - Mail to: WVU Pediatric Endocrinology PO Box9214 Morgantown, WV 26506

4. Allow 5-7 days for our office to receive and review completed forms. We will then send completed forms with a physician's signature to your child's school using the information provided on the form. If a verified fax number or e-mail is not included on the Diabetes School Orders form, this will prolong the process.

Please Note: Schools will not permit children to ATTEND school without the Diabetes School Form being completed and signed by a physician. If you have any questions, please contact our office.