

HOST LIST OF AVIAN BROOD PARASITES - 1 - ANATIDAE

Peter E. Lowther, *Field Museum*

version 12 Oct 2023 (original list 19 Oct 2009, illustrations added 08 Jun 2012; new material added 29 Apr 2013, 18 Jul 2017)

Heteronetta

Heteronetta Salvadori 1865, Atti. Societa Italiana di Scienze Naturale e Museo Civico di Storia Naturale, 8, p. 374.

Black-headed Duck, *Heteronetta atricapilla* (Merrem) 1841, Allgemeine Encyclopadie der Wissenschaften und Kunste in alphabetischer Folge von genannten Schriftstellern bearbeitet und hrsg. von J. S. Ersch und J. G. Gruber, sect. 1, 35, p. 26.



General life history information given by Weller (1968, Lyon and Eadie 2013; see also Lowther 2010). Host list from Weller (1968), Höhn (1975), Lyon and Eadie 2013, Cabrera et al. 2017; see also Rees and Hillgarth (1984). Livzey (1986) presents waterfowl phylogeny showing Black-headed Duck to be sister-group to stiff-tails.

Species known to have reared this brood parasite indicated by “%”. In captivity, reported hosts have included Rosy-billed Pochard *Netta peposaca* (Powell 1979, Rees and Hillgarth 1984), Red Shoveler *Anas platalea* (Todd 1979¹), and Common Moorhen *Gallinula chloropus* and Silver Teal *Anas versicolor* (Powell 1979². Rees and Hillgarth 1984).

The Maguari Stork *Ciconia maguari* had been listed originally as a host based on supposed parasitic eggs which were later identified as being those of Rosy-billed Pochard *Netta peposaca* (see Weller 1968); subsequent records returns this species to this list of hosts (Cabrera et al. 2017).



ANHIMIDAE

Southern Screamer, *Chauna torquata*

ANATIDAE

Fulvous Tree-Duck, *Dendrocygna bicolor*

Black-necked Swan, *Cygnus melanocoryphus*

Coscoroba Swan, *Coscoroba coscoroba*

White-cheeked Pintail, *Anas bahamensis*

% **Rosy-billed Pochard, *Netta peposaca***

PODICIPEDIDAE

Great Grebe, *Podiceps major*

ARAMINIDAE

Limpkin, *Aramus guarauna*

RALLIDAE

Spotted Rail, *Paradirallus maculatus*

Plumbeous Rail, *Paradirallus sanguinolentus*

% **Red-fronted Coot, *Fulica rufifrons***

% **Red-gartered Coot, *Fulica armillata***

White-winged Coot, *Fulica leucoptera*

RECURVIROSTRIDAE

Black-necked Stilt, *Himantopus mexicanus*

LARIDAE
 Brown-hooded Gull, *Chroicocephalus maculipennis*

CICONIIDAE
 Maguari Stork, *Ciconia maguari*

ARDEIDAE
 Black-crowned Night Heron, *Nycticorax nycticorax*

THRESKIORNITHIDAE
 White-faced Ibis, *Plegadis chihi*
 Bare-faced Ibis, *Phimosus infuscatus*
 Roseate Spoonbill, *Ajaia ajaia*

ACCIPITRIDAE
 Snail Kite, *Rostrhamus sociabilis*
 Long-winged Harrier, *Circus buffoni*³

FALCONIDAE
 Chimango Caracara, *Milvago chimango*

Acknowledgments. Illustration and map taken from Neotropical Birds Online; photo of bird at Rio Grande do Sul, Brazil, 22 November 2009 © Cláudio Dias Timm; map provided by Robert S. Ridgely.

References:

- Cabrera, M. B., D. Montalti and L. N. Segura. 2017. Breeding phenology and new host list of the Black-headed Duck (*Heteronetta atricapilla*) in Argentina. *Wilson Journal of Ornithology* 129: 311-316.
- Hellmayr, C. E., and B. Conover. 1948. Catalogue of birds of the Americas and the adjacent islands in Field Museum of Natural History including all species and subspecies known to occur in North America, Mexico, Central America, South America, the West Indies, and islands of the Caribbean Sea, the Galapagos Archipelago, and other islands which may be included on account of their faunal affinities. Part 1, no. 2. Spheniscidae-Gaviidae-Colymbidae-Diomedidae-Procellariidae-Hydrobatidae-Pelicanoididae-Phaethontidae-Pelecanidae-Sulidae-Palacrocoracidae-Anhingidae-Fregatidae-Ardeidae-Cochleariidae-Ciconiidae-Threskiornithidae-Phoenicopteridae-Anhimidae-Anatidae. Field Museum of Natural History, Zoology Series, vol. 13, Publ. No. 615.
- Höhn, E. O. 1975. Notes on Black-headed Ducks, Painted Snipe, and Spotted Tinamous. *Auk* 92: 566-575.
- Livzey, B. C. 1986. A phylogenetic analysis of Recent anseriform genera using morphological characters. *Auk* 103: 737-754.
- Lowther, P. E. 2010. Black-headed Duck (*Heteronetta atricapilla*), Neotropical Birds Online (T. S. Schulenberg, ed.). Cornell Laboratory of Ornithology, Ithaca, NY; retrieved from: http://neotropical.birds.cornell.edu/portal/species/overview?p_p_spp=21158;
account currently available in Birds of the World as:
 Lowther, P. E. 2020. Black-headed Duck (*Heteronetta atricapilla*), version 1.0. In *Birds of the World* (T. S. Schulenberg, ed.). Cornell Lab of Ornithology, Ithaca, NY, <https://doi.org/10.2173/bow.blhduc1.01>
- Lyon, B. E., and J. M. Eadie. 2013. Patterns of host use by a precocial obligate brood parasite, the Black-headed Duck: ecological and evolutionary considerations. *Chinese Birds* 4: 71-85.
- Powell, A. 1979. Cuckoo in the nest. *Wildfowl News* 81: 15-16.
- Rees, E. C., and N. Hillgarth. 1984. The breeding biology of captive Black-headed Ducks and the behavior of their young. *Condor* 86: 242-250.
- Todd, F. S. 1979. *Waterfowl. Ducks, geese and swans of the world*. Seaworld Press, San Diego, CA.
- Weller, M. W. 1968. The breeding biology of the parasitic Black-headed Duck. *Living Bird* 7: 169-208.

Notes.

1. Todd (1979: 314): "Breeding success was achieved for the first time in 1977 at the Wildfowl Trust, and the Assistant Director, Mike Lubbock, has made a number of observations that have added considerably to our knowledge of the breeding biology of this interesting species. In one case, a pair of black-headed ducks was observed to *force* an incubating Argentine red shoveler from the nest, and within five minutes the female *Heteronetta* was seen to lay an egg of her own."
2. Based on published information, the identity of this teal host is not certain. Captive breeding as described by Powell (1979: 16) indicated 3 different host species: "Between the 9th and 18th of May [1979], six more Black-headed Duck eggs have been laid: two in the same nest [of Rosy-billed Pochard], two in a Moorhen nest and two in a Cinnamon Teal nest and , ... " ; but the host environment for these Black-headed Ducks at the Wildfowl Trust has been described slightly differently in terms of potential host species available.

Rees and Hillgarth (1984: 242-243) described captive conditions as: "In 1974 the Wildfowl Trust collected eggs of Black-headed Ducks in Argentina in order to establish a captive breeding population of this species at Slimbridge, Gloucestershire, England. The five males and two females from these eggs were pinioned and placed in a pen (36 x 19 m) in which the terrain was designed to resemble their natural habitat (M. R. Lubbock, pers. comm.). The pen contained a pond (380 m² surface area) and four small islands. Rushes and sedges were planted at the water's edge so that the area was like a miniature marsh. **Other South American birds** [emphasis added] were put in the enclosure to provide suitable hosts for the Black-headed Ducks. These included Rosy-billed Pochards, Lake Ducks (*Oxyurua vittata*), Southern Versicolor Teal (*Anas versicolor fretensis*), Red Shovelers (*A. platalea*), and Black-necked Swans (*Cygnus melanocoryphus*). Wild Mallards (*A. platyrhynchos*), Gadwalls (*A. strepera*), and Moorhens (*Gallinula chloropus*) also bred in the pen. The first nine captive bred Black-headed Ducks were reared at Slimbridge in 1977."

Powell (1979: 15) describes these same conditions in the Wildfowl Trust's newsletter as: "At Slimbridge where the Black-headed Duck first bred in captivity in 1977, (an account appeared in 'Wildfowl News' No. 77) six pairs are at present kept in a pen with three pairs of Rosybill [= Rosy-billed Pochard, *Netta peposaca*], three pairs of Argentine Ruddy Duck [= most likely Lake Duck, *Oxyura vittata*; if only truly Argentinean species shared the pen; Ruddy Duck, *Oxyura jamaicensis* does not naturally occur in Argentina], one pair of Cinnamon Teal [= *Anas cyanoptera*; or possibly – or more likely given the species listing in Rees and Hillgarth 1984, Silver Teal, *Anas versicolor*] and one pair of Black-necked Swans [*Cygnus melancoryphus*] -- all **natives of Argentina** [emphasis added] and all ground nesting species."
3. Nest may have been abandoned when parasitized.