

Capsulectomy

Pre & Post Operative Instructions

Understanding Your Surgery with General Anesthesia:

- **General Anesthesia:** This type of anesthesia renders you completely unconscious during the procedure, ensuring you feel no pain and have no memory of the surgery. It is typically used for more extensive operations and requires careful monitoring by an anesthesiologist.
 - **Instructions:** Nothing to eat or drink 8 hours before surgery. This includes water, mints, candy, or gum.
 - **Post-Surgery Transportation:** It is mandatory to arrange for a ride home from surgery with a family member or friend. Patients may not use taxis or Ubers.
 - **Why?:** After surgery, especially with general anesthesia, it's important for your safety to have someone responsible to take you home and provide care for at least 1-2 days. This ensures that you have the necessary support during the initial critical recovery period. If you do not arrange for transportation and care for at least 1-2 days, then your surgery may be canceled, at your expense.

Shopping List

- Tylenol (Acetaminophen): Alternative to narcotic pain medication if pain is not severe. Do NOT take NSAIDS (such as Ibuprofen & others).

Pre-Operative Instructions

Medication Management Before Surgery: Guidelines for Minimizing Bleeding Risks

These are general recommendations. For more information please refer to the email: Preoperative Instructions for Your Upcoming Surgery &/or contact our office for a safe plan tailored to your health needs and surgical procedure

- **Aspirin and NSAIDs:** Please discontinue use at least 3 weeks before your surgery, unless otherwise directed by your surgeon.
- **Anticoagulants (Warfarin, Heparin, Direct Oral Anticoagulants):** The discontinuation schedule varies. Please consult your surgeon at least 3 weeks before surgery to discuss a tailored plan, which may include temporary adjustments to your regimen or bridging therapy.
- **Herbal Supplements and Vitamins:** Discontinue any herbal supplements and vitamins that can increase bleeding risk, such as vitamin E, fish oil, ginkgo biloba, and garlic, at least 2 weeks prior to surgery.
- **ADD / ADHD medications** are to be avoided the day before surgery.
- **Contrave (Naltrexone-containing meds):** Patient must consult with the prescribing physician about when & how to stop this medication prior to surgery.
- **Weight Loss Meds (Ozempic, Wegovy, Semaglutide, ect.), & Biotin.:** are to be stopped 2 weeks prior to surgery.
- **Illicit Drug Use:** Please be honest about any illicit drug use during the preoperative assessment. Your safety is our priority and accurate medical information is crucial for planning a safe and effective surgical procedure.
 - Illicit drug use can have significant interactions with anesthesia and other medications administered during surgery, potentially leading to complications.
 - **Examples of the most important substances to mention include:**
 - Cocaine & Meth.

Consultation Is Key: If you are taking any of the above medications or supplements for chronic conditions, do not stop them without first consulting with the physician who prescribed them and your surgeon.

Together, we will develop a safe plan tailored to your health needs and surgical procedure.

4 Weeks Before Surgery

- **Reducing/Stopping Nicotine:** We strongly recommend reducing or ideally stopping nicotine (including cigarettes, oral tobacco, e-cigarettes with nicotine, patches, lozenges, and/or gums) for 4 weeks before and after your surgery.
 - **Why?:** Nicotine can significantly hinder your healing process and increase the risk of complications. Nicotine constricts blood vessels, which can impair blood flow to the surgical area, affecting healing and recovery. Any reduction in nicotine exposure is beneficial and contributes positively to your surgical outcome.

2 Weeks Before Surgery

- **Alcohol Consumption:** We encourage patients to reduce or stop alcohol consumption for at least 2 weeks before and after surgery.
 - **Why?:** Alcohol can increase the risk of bruising, swelling, and can prolong the healing process. Reducing alcohol intake, even if complete cessation is challenging, but can significantly improve your recovery experience.

The Day of Surgery: Your Preparation Guide

On the day of your surgery at Asia Pacific Surgery, we prioritize your comfort, safety, and the success of your procedure. To help everything go smoothly, please follow these guidelines carefully:

- **Shower or Bath:** On the day of surgery, please take a shower or bath using soap (either antimicrobial or non-antimicrobial).
 - **Avoid Applying Products:** Do not apply any lotions, perfumes, or other products to your skin after bathing.
- **Medications:**
 - **Blood Pressure Medications:**
 - Take your blood pressure medication as usual, including the day of surgery. On the day of surgery, take it with only a small sip of water.
 - **Diabetes Medications:**
 - DO NOT take any oral and/or injectable diabetic meds the night before or the morning of surgery.
 - Only take medications approved or prescribed by your surgeon with a small sip of water.
- **Fasting:** Adhere to the fasting instructions provided by your surgical team. Typically, this means no eating or drinking, including water, for 8 hours before your surgery. Check your paperwork or email our office for further guidance, if needed. This information will be on the 1st page of this document.
- **Jewelry and Valuables:** Remove all jewelry and refrain from bringing valuables to the clinic, including piercings from the neck up. We cannot take responsibility for lost items.
- **Makeup and Skincare Products:** DO NOT wear any makeup or apply skincare products on the day of your surgery.
- **Menstruation and Surgery**
 - If you are on your menstrual period on the day of surgery, you may wear a sanitary pad or disposable underwear.

- For safety reasons, internal menstrual products (such as tampons or menstrual cups) are not used.
- After surgery, if you need menstrual supplies before leaving the facility, pads can be provided upon request. Application of any menstrual product is performed by the patient.
- **Clothing:** Opt for loose, comfortable clothing that doesn't need to be pulled over your head. A button-up or zip-front top is ideal for after the surgery.
- **Communicating Questions or Concerns:** If you have any last-minute questions or concerns, please get in touch with us immediately. See the last page of this document for contact information.

Post-Operative Instructions

Medications

1. Tylenol (Acetaminophen):
 - Pain Relief: Tylenol is safe to take for managing pain. Please avoid NSAIDs such as Ibuprofen and Aleve, as they can increase the risk of bleeding.
2. Narcotic Pain Medication:
 - Prescription Medication: If you received a prescription for narcotic pain relievers, take them as directed for pain management. Be aware that these medications can cause constipation.
 - Preventing Constipation: You will be prescribed a laxative and a stool softener to prevent constipation. It's important to start taking these with your first dose of narcotic pain medication. Additionally, increasing dietary fiber, fluid intake, and physical activity can help manage constipation.
3. Zofran (Ondansetron):
 - Nausea Prevention: Zofran is prescribed to help prevent nausea and vomiting, which can be common after surgery. Take this medication as directed, particularly before meals or as soon as you feel nauseated.
4. Antibiotics:
 - Infection Prevention: If you've been prescribed antibiotics, it's crucial to complete the full course as directed, even if you start feeling better. This helps prevent infection and ensures the effectiveness of the treatment.
5. Sinnech (Arnica Montana):
 - Reducing Swelling and Bruising: Sinnech is a homeopathic Arnica montana regimen that you'll take to help reduce swelling and bruising. This can potentially speed up your recovery process. Please follow the specific dosage instructions provided.

Nutrition

- Hydration: Stay hydrated with plenty of water. A general guideline is to aim for at least 8-10 glasses of water per day (about 2 to 2.5 liters). Limit or avoid caffeine and alcohol.
- Meal Approach: If you have a reduced appetite, eat small, frequent meals.

Activity

- **Movement is Important:** Make sure you are up and walking around immediately after your surgery. This helps prevent Deep Vein Thrombosis (DVT), a condition where blood clots form in the veins, usually in the legs. Regular movement improves blood circulation and reduces the risk of clots. Start with gentle walks around your home and gradually increase activity as recommended by your surgeon.
- **Sleep:** Sleep on your back with your head elevated about 30-40 degrees (2-3 pillows). Do not sleep on your side. Keeping your body more upright will minimize swelling. Continue this for one week.
- **Exercise:** Light physical activity may be resumed 2-3 weeks after surgery. Remember to start easy and build back up to your previous exercise levels. At 6-8 weeks or when further instructed by your surgeon, more intense exercise can be started. Just know that swelling may transiently be worse with exercise.
- **Arm Movement:** Limit your arm use to daily activities such as brushing your teeth, eating, and combing/shampooing your hair. Avoid vigorous movement and do not lift more than 5 lbs. Let pain be your limiting factor. If it hurts, please do not do it.
- **Return to work:** Most patients require approximately 5-7 days off work depending on their job responsibilities. Returning to work with a light schedule initially or even part-time can be beneficial as well.
- **Driving:**
 - Do not drive while taking narcotic pain medication.
 - Most patients can resume driving safely after 1–2 weeks once they are off narcotics, have good arm mobility, and feel safe controlling the car.
 - Avoid long drives or manual-shift driving for about 3 weeks.

- **Sexual Intercourse:** Sexual activity can be resumed when you feel ready with no restrictions. Incisions should be well healed. Patients typically feel ready after one month. Please be sure to keep the following in mind:
 - Do not touch the breasts
 - Do not lay on your stomach or on your side
- **Sleeping:** It is advised for the first few weeks to sleep on your back to avoid putting too much pressure on the breasts. Do not lie on your chest.

Note: These are general guidelines and may vary depending on your individual recovery, lifestyle, and specific circumstances. Please follow your surgeon's personalized advice to ensure a safe and effective recovery.

Your Recovery Bra

You will be provided with a supportive surgical bra after surgery.

- The bra should feel supportive but not constricting or overly compressive.
- Wear it 24 hours a day for the first month, with two short breaks daily.
- To make daily wear easier, we recommend purchasing two bras so you can alternate as needed while keeping one clean.
- You may purchase your own garment if you prefer, as long as it is:
 - Made of a soft material
 - Opens and closes in the front (no garments that go over the head or close in the back)
 - Contains no underwire

Wearing the garment as directed is an important part of your recovery and helps support optimal healing.

Bathing & Swimming

- Showering: You may shower, with assistance, 24–48 hours after surgery. Your incisions are covered with dressings, which can be removed 24 hours after your surgery.
- Hot Tubs/Baths/Swimming Pools. No tub baths or Jacuzzi until your incisions have healed, and approved by your surgeon, which is usually around 2 weeks. It is best to wait one month for hot tubs as they tend to have more bacteria than regular chlorinated swimming pools.

How to Care for Your Incisions

- Incisions: Your incisions are covered with a dressing, which can be removed 24 hours after your surgery.
- Drainage Tubes: Drainage tubes may be necessary for a capsulotomy or capsulectomy surgery.
 - Please see the [JP Drain Home Care Instructions](#) document provided to you for more information.
- Stitches: All stitches are dissolvable.
- Surgical Glue: In some cases, surgical glue may be used to seal the incisions. This glue will naturally fall off over time and does not require any special care.

What to Expect

- Bruising: You can expect to have some bruising. Most bruises will resolve after about 2-3 weeks. The bruise will go from a purplish color to a yellow/green shade as it starts to resolve.
- Swelling: You can expect swelling particularly in the top half of the breasts. This can take 3-4 weeks to improve and the swelling is responsible for the feeling that the implants are high. As the swelling resolves, it seems as though the implants drop into a lower position.
- Itching: Itching at the incision sites is normal for a few days. You may take Benadryl to help with this.
- Pain: It is normal to experience tightness, pressure, soreness, itchiness, and fatigue for several days to weeks following surgery as your skin and muscles adjust to your breast implants.

- **Sensory Changes in Skin:** You may feel reduced or heightened sensation in the nipples, incision sites, and breast tissue, which is normal. You can expect a return of normal sensation after a few weeks to months.
- **Crackles/Sloshing Sounds:** Air and fluid are left behind as a normal part of the surgery and it takes time for it to be resorbed by your body. These sounds are normal after a breast augmentation and will resolve in 5-7 days.

Emotional Recovery

It's normal to experience a range of emotions after surgery, especially when the appearance of your body changes. Some patients feel relief, while others may go through moments of doubt or grief. Please know that these feelings are valid and part of the healing process. You're not alone, and support is available if you need it.

Signs of Infection or Complications

It's important to monitor your recovery and be aware of any signs that may indicate an infection or complication. Contact your surgeon immediately if you experience any of the following:

- Increased redness, warmth, or swelling around the incision site
- Pus-like or foul-smelling drainage from the wound
- Fever over 101°F (38.3°C)
- Severe pain that is not relieved by prescribed medication
- Excessive bleeding or new bruising
- Persistent chills or feeling unwell

Prompt communication with your healthcare team ensures the best possible recovery. Do not hesitate to reach out if you have concerns.

Follow-up Care

- **Prioritizing Appointments:** Attending follow-up appointments is crucial for monitoring your recovery & ensuring the effectiveness of your treatment.

Patient Communication Guide: Who to Contact Post-Operation

Type of Concern	Contact Information	Instructions/Examples
Non-Urgent Questions or Concerns for Dr. Ching or Dr. Long	Email: clinicalrn@drshimching.com	<ul style="list-style-type: none"> • General recovery questions • Appointment inquiries • Prescription refills
Urgent Questions or Concerns	Office Phone: (808) 585-8855 <u>After 5pm or on weekends:</u> Dr. Ching & Dr. Long @ Physician's Exchange (808) 524-2575	<ul style="list-style-type: none"> • Swelling, bruising, or redness • Severe pain • High fever • Unusual discharge or odor • Uncontrolled bleeding
Medical Emergencies	Call 911 or go to the nearest emergency room	<ul style="list-style-type: none"> • Difficulty breathing • Chest pain • Severe calf pain • Signs of heart attack or pulmonary embolism

Our Commitment to You:

We are here to support you throughout your recovery journey. Your safety is our priority, and we encourage open communication. By guiding you on the most appropriate communication channels for your concerns, we aim to provide efficient and effective care, ensuring you feel fully supported & cared for.