



INGREDIENT HIGHLIGHTS

- Niacinamide (Vitamin B3), Hyaluronic Acid (Sodium Hyaluronate), Methyl Glucoside Phosphate Proline Lysine Copper Complex, Sodium Chondroitin Sulfate, Glycosaminoglycans, Pea Extract, Glucosamine HCl, Panthenol (Provitamin B5), Bambusa Vulgaris Leaf/Stem Extract

INGREDIENTS

- Water, Glycerin, Niacinamide, Sodium Chondroitin Sulfate, Methyl Glucoside Phosphate Proline Lysine Copper Complex, Glycosaminoglycans, Bambusa Vulgaris Leaf/Stem Extract, Pisum Sativum (Pea) Extract, Glucosamine HCl, Sodium Hyaluronate, Panthenol, Phenoxyethanol, Ethylhexylglycerin, Polysorbate 20, Disodium EDTA

PLUMP

DESCRIPTION



Employing multiple revolutionary active ingredients, PLUMP is clinically proven to deliver equivalent results to one injection of collagen filler in just two weeks of regular use.

The combination of copper, proline, lysine, hyaluronic acid, and niacinimide triggers the production of energy in the mitochondria of the senescent fibroblast, safely reactivating collagen I, collagen III, and elastin production. Known as the "youth collagen," collagen III presence naturally decreases starting at age 8 or 9, and is virtually impossible to reactivate. Golden Bamboo Extract and Pea Extract deliver an 85% increase in hyaluronic acid synthesis during an in vitro test, plus a decrease in skin laxity of 50% after four weeks. The result is a painless, topical alternative to injectable fillers and toxins.

In addition to building collagen, Needle Free is fortified with glycosaminoglycans which preserve and protect the structure of the ECM complex from physical forces like pollution, UV rays and aging; leading to an improvement in the skin's barrier function and a reduction in the appearance of rosacea over a 12 week period.

BENEFITS

- Reduces wrinkle appearance and volume by 15% in only 2 weeks
- Quickly delivers equivalent results to one injection of collagen filler
- Firmness increased by 13 times in 15 days
- A painless, topical alternative to injectable fillers and toxins
- 190% Increase in elastin production after 11 days
- 179% Increase in collagen I production , and 194% Increase in collagen III production after 11 days
- 85% increase in hyaluronic acid synthesis
- 50% decrease in skin laxity after 28 days
- Improvements sustained after two months of treatment
- 80% inhibition of MMP enzyme activity (types 1 and 2)
- Improves the skin's barrier function and shows a reduction in the appearance of rosacea over a 12 week period

DIRECTIONS FOR USE

- Apply 4 to 5 drops onto fingers and smooth over face and neck, avoiding eye area. Allow to absorb before applying additional products. Can be used twice a day or as directed by physician.