

## After Breast Lift Instructions

You will have a bra in place after surgery. This should remain in place for the first 24 hours after surgery.

It is very important that you **do not smoke, do not wear a nicotine patch, or do not chew nicotine gum** in the first 3 weeks after surgery. Severe scarring may result from failure to observe this precaution.

**Medications** (you may be given all or some of the following medications depending on your surgery):

- Percocet: You may take it every 4 hours as needed. Do not drive a vehicle or operate heavy machinery while using this medication.
- Zofran: In the event that you experience nausea or vomiting, take this medicine. While you may not need it, it is advisable to fill the prescription just in case.
- Colace: Your pain medicine can cause constipation, and excess strain and pressure can not only be unpleasant, but may hurt your incisions. Please take this while on the Pain Medication.
- Zanaflex: This will help relax the muscles involved in your procedure, which will also help with pain control.

Although you have been given a prescription for pain medication, you may only need regular Tylenol or Ibuprofen and should try it first. If your stomach becomes irritated or upset, stop the medication. **Do not use aspirin containing products which may increase your chances of bleeding complications.**

**Surgical site care:**

- Leave bandages in place until tomorrow.
- Change bandages tomorrow.
  - Remove gauze, shower gently and pat dry.
  - Do not allow direct high pressure shower spray to hit incisions.
  - Do not soak incisions.
  - Leave steristrips in place.
- Use fluffed gauze to support breasts inside bra to push them toward the middle (fluffs go along outer edge of each breast.)
- Wear bra 23 hours a day for one month (to support the breasts).

**Activities:**

- Sleep with head of bed elevated 30 degrees.
- **No heavy lifting for two (2) weeks.**
- Ambulate today/ tonight.
- 20 pound lifting limit.

- No running or impact sports for 1 month after your procedure.
- Do not engage in sexual activity for a minimum of two weeks.

#### **Returning to Particular Activities**

- You may return to nonimpact aerobics at 2 weeks after surgery (stair master, low-speed elliptical).
- You may begin running at 4 weeks after surgery, but wear a sports bra over your surgical bra for the next 2 weeks (post-op week 5 and 6).
- You may begin push-ups, pull ups and swimming 6 weeks after your procedure.
- With all activities, it is important to listen to your body. Don't push too hard, take it "low and slow."
- Avoid quick, hard movements that could break a suture or tear the incision apart. Avoid excessive lifting that can cause increased pressure and possible bleeding.

#### **Diet:**

- As tolerated, No restrictions.
- We would recommend you start resuming your normal diet gradually. Start with thin or thick liquids and advance to a general diet as tolerated.
- Drink plenty of water (eight 8-oz glasses each day).
- Do not drink any alcoholic beverages for 3 days. Never drink alcoholic beverages while on prescribed pain medication.

#### **Follow-up appointments:**

- 2 weeks after surgery: Your steristrips will be removed at this time. After your steristrips are removed you may use regular moisturizer daily. We will discuss scar care at this visit. Be sure to apply sunscreen to your suture lines to prevent pigmentation issues.
- 2 months after surgery: Monitor healing and swelling.
- 6 months: Final check to make sure everything is going great after your breast lift procedure.

#### **Watch for the following signs and symptoms, and notify the office immediately if these occur:**

- Fever or blistering.
- Redness that looks like sunburn, tenderness or warmth in the area of redness.
- Sudden increase in one-sided pain.
- Swelling that is much greater on one side.
- Bruising is **NORMAL**.

Please call the office if you have any questions.