

Hair Transplant Instructions

Before your procedure

- For 14 days prior to your procedure, do not take aspirin, naproxen, ibuprofen, Omega 3, Fish Oil, Vitamin E, ginkgo biloba, or any multivitamins that contain Vitamin E. Many medications contain aspirin, so please be sure to read all medication labels. You may take acetaminophen (e.g. Tylenol) in place of aspirin.
- For 3 days prior to your procedure avoid drinking alcohol and do not get a sunburn.
- **Start taking Keflex (antibiotic) the day before your procedure. Continue taking until complete.**
- **Avoid caffeine or alcohol the night before and the morning of your procedure.**
- Make arrangements for a driver to drive you to and from your procedure.
- If you have gray hair and you color it, make sure it gets colored within 3 days prior to your procedure to permit better visualization of the hairs. You will be able to color it again 6-8 weeks after the procedure. A semi-permanent color should be used instead of a permanent hair color that may cause damage to your new grafts. Permanent hair color can be used 3 months after your surgery.
- For best healing, please refrain from smoking for 1 week prior to and 1 week after your procedure.
- Vitamin C (1000 to 2000 mg a day) for 1 week prior to your procedure may also help with healing.
- Wash your hair the day before surgery with baby shampoo.
- Please consider what hairstyle you will wear for your procedure. The donor site will need to be shaven down to skin level. A short haircut will allow others to see you have had surgery vs. a long haircut will enable us to comb your hair over the donor site and provide some camouflage.

The day of your procedure

- Have a regular breakfast, but **no caffeine in the morning**. You will be provided lunch and offered snack breaks during your procedure.
- Any prescribed medicines (e.g., blood pressure) should be taken the morning of your procedure.
- Wear a shirt that can be buttoned in the front rather than a pull-over. Remove all jewelry, watches, etc.
- Be prepared to remove contact lenses before your procedure if you normally remove them to sleep.
- Please feel free to bring your iPod or iPad to use to during your procedure.
- Friends and family are not allowed in the surgery room, but may stay in the waiting room.

After your procedure

- **Medications:** These are the medications that will be prescribed two weeks prior to your procedure. You will need to pick up the written scripts at our office and have them filled at the pharmacy of your choice; bring them with you on the day of your procedure. If you are not able to take any of them (i.e. due to allergies) please let us know before your procedure.
 - Norco: This is a pain medication. Take 1 tab by mouth every 4 hours as needed for pain. Please bring this prescription with you to the office the day of your procedure.
 - Valium: This is an anti-anxiety medication. Take 1 tab by mouth every 8 hours as needed for anxiety or itching. Please bring this prescription with you to the office the day of your procedure.
 - Keflex: This is an antibiotic. Take 1 capsule by mouth 4 times daily starting the day before your procedure. Continue taking until completed.
- Alcoholic beverages, ibuprofen and aspirin are not to be taken for 48 hours after surgery. You may transition from the prescribed pain medication to extra strength Tylenol after your procedure.

24-48 Hours Post-Op

- AVOID CONTACT TO DONOR SITE AND NEWLY IMPLANTED GRAFTS.
- Do not wash your hair. **Use Evian Spray Water** to keep the grafts moisturized. We recommend sleeping in a shower cap to help keep the grafts moisturized during the night.
- You may also use **Hair Medica Copper K Calming Spray** 2-3 times daily for additional moisturization to the scalp after your procedure. Use as directed for the first 72 hours after your procedure, then as needed for the next 2 weeks.
- Avoid any strenuous physical activity for the first 7-10 days to prevent loss of grafts or increased swelling. Swimming may be resumed 14 days after your procedure.
- No direct sun exposure to the scalp for more than 30 minutes at a time for the first 5 days. Wearing a hat provides adequate protection. We recommend wearing an adjustable (not fitted) cap. Do not wear a hat that collapses or touches your scalp (beanie, wool cap, etc.). The cap should not come in contact with the newly transplanted grafts. If you plan to be outside for more than 30 minutes, you may use a spray sunscreen after 2 weeks.
- A pressure dressing will be applied to the back of the scalp where grafts were harvested. This dressing should remain in place for 24 hours. After the pressure dressing is removed, use **Copper K Calming Spray** over the donor area twice daily until healed.

For Sleep:

- **Frontal Hairline Restoration:** it is recommended that you sleep in a flat position to move fluid back from eyes and prevent swelling.
Crown Restoration: you may sleep slightly elevated.

For Minor Bleeding:

- Occasionally there is some slight bleeding from the back (donor area) and or several graft sites the first 24 hours. Simply apply gentle but **firm** pressure with a clean damp cloth or gauze for 5-10 minutes until the bleeding stops. **Press only. Do not rub.**

48 Hours Post-Op

- **Do not wash your hair for 48 hrs. after surgery.**
- **You may gentle cleanse your hair 48 hrs. after surgery as follows.**
 1. Wet your head in the shower with warm water, using a cup or pitcher to pour it over the scalp (top, sides, and back).
 2. You may use a gentle shampoo (Baby Shampoo) and lukewarm (NOT HOT) water. Place Shampoo and water in a cup. Create a lather and pour this mixture onto your entire scalp and spread the shampoo over the sides and back of the scalp. You may gently massage the mixture with your fingers **only on the donor site. It is important that you do not touch the recipient areas with your fingers as this may cause the grafts to dislodge.**
 3. Gently rinse off the shampoo, then air dry. Do not allow full strength streaming water from the shower head to make contact with your scalp. This may cause your grafts to dislodge, especially within the first 5 days.
 4. Wash your scalp daily for the first 2 weeks.
 5. Continue using water to spritz the grafts and keep them moisturized.
- You may wash your hair like normal 2 weeks after your procedure.
- Combing, performed gently, is permitted on the third day. When combing and styling your hair, make sure to do so while the hair is wet. Comb the hair in an “up and over” manner. This will eliminate the chance of scratching the scalp and dislodging the grafts. Do not use any hair styling products for the first 14 days.
- You may resume low-level laser therapy (Capillus Hair Caps) **48 hours** after your procedure.

72 Hours Post-Op

- If swelling occurs, it will usually appear on the second or third day after surgery. Swelling is part of the healing process and will usually disappear within 5-7 days.
- You may also use an ice pack or bag of frozen peas over the eyebrows for 15 minutes every hour starting immediately after surgery and continuing for the next 72 hours. **Do not place ice over the grafts.**

What to expect

- You may have numbness at the graft and donor sites. This is normal. It can take several weeks, occasionally several months to resolve.
- Scabbing will form over the transplant sites and solidify the first few hours after the procedure. These scabs will naturally fall off within 2 weeks. Some of the transplanted hairs may fall off with the crusts. Do Not Panic for these are only the hair shafts, while the hair follicles remain in the scalp. **Do not pick or scratch at the scabs at any time. You can endanger the healing of the graft and could cause scarring or infection.**
- In as soon as 3 to 4 months, the new hair shafts will begin to grow in a natural fashion and will continue to grow at a normal rate of one half to 1 inch a month. Sometimes it can take 7 to 9 months or more for all the hairs to start growing. Be patient.
- To accelerate the growth of the new hairs we recommend patients begin applying 5% minoxidil once or twice a day starting 1 month after the procedure. Female patients should use 2% minoxidil. If the minoxidil should cause redness or irritation of the scalp, stop using it for 1 week then resume once every other day. Once the hairs are growing in, at around 4 months, the minoxidil can be stopped, and the transplanted hairs will continue to grow. The occasional patient is sensitive to minoxidil (scalp itching, chest palpitations, lightheadedness) in which case it should be stopped.
- Propecia can continue to be used.
- Be Patient for the hairs to grow in. As they emerge from the scalp, occasionally there will be a few acne-like pimples. If this happens, apply warm soaks and antibiotic ointment several times a day for 3 days. If not resolved, please call the office.
- You may have your hair cut as soon as 3 weeks after the procedure. Hair coloring is permitted 6-8 weeks after your procedure. A semi-permanent color should be used instead of a permanent hair color that may cause damage to your new grafts. Permanent hair color can be used 3 months after your surgery.

We are here to make this process as smooth and comfortable as possible for you.

Watch for the following signs and symptoms. Notify the office immediately if these occur:

- Fever or blistering.
- Excessive bleeding that cannot be stopped by firm pressure and application of ice to the area for a period greater than 20 minutes.
- Minor bleeding is **NORMAL**.